Global comparisons of total O3 columns from SCIAMACHY, GOME and OMI

A. Bracher, L. N. Lamsal, M. Weber, M. Coldewey-Egbers, J. P. Burrows
University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de

Introduction

Global stratospheric ozone columns derived from UV nadir spectra measured by OMI, SCIAMACHY and GOME are compared. Data sets for SCIAMACHY are the products from the weighting function DOAS algorithm (WFDOAS Version 1; see Coldewey-Egbers et al. 2005) and the most recent ESA operational product versions 5.04/5.01, GOME WFDOAS and OMI level-3 data are taken from the are from IUP webpage and TOMS website, respectively Because of different spatial resolutions (GOME 320x40, SCIAMACHY 60x30, OMI 24x18 km), data from 2003 and 2005 are binned into appropriate spatial bins and then compared.

GOME Weighting Function DOAS (WFDOAS): Algorithm and Validation

Main feature of WFDOAS (see Coldewey-Egbers et al. (2005):

- Taylor expansion of optical depth spectrum around climatological total ozone (TOZ_{clim})
- Modelled radiative transfer quantities: reference OD and weighting function parameterised by : TOZ, eff. albedo, eff. scene height, viewing geometry
- Improved Raman correction
- Ozone temperature correction
- Cloud cover and effective scene height from oxygen A-band absorption (Koellemeijer et al. 2001) including ghost vertical column correction
- Effective UV albedo (surface reflectance) from 377nm
- Spectral fitting window: 326.6-335nm
- Climatological ozone from TOMS V7 profile shape climatology (Wellemeyer et al. 1997)

Comparison of GOME WFDOAS with Brewer data (see Weber et al. 2005)

Well-maintained Brewer data- good source for satellite O3 validation since Brewer has negligible O3-T dependence in their wavelengths



GOME WFD compared with Brewer data: within 1% and negligible seasonal cycle!

Comparisons of GOME WFD, SCIAMACHY 5.01/5.04 and SCIAMACHY WFD (from Bracher et al. 2005)

SCIAMACHY (=SCIA) WFD V1, SCIA operational product 5.04 and GOME WFD V1 are compared at several days during Jan to June 2003, the comparison between SCIA op. 5.04 and GOME WFD V1 was performed for the whole data set during this time frame. GOME is a good candidate for validation of SCIA: equator crossing time 10:30 LT (SCIA 10 LT) and same orbit. Data sets are binned into 2.5° lat x 2.5° lon.

Global tO_3 distributions



Overview

SCIA WFD to GOME WFD 12 May 2003 | **SCIA WFD to SCIA 5.04 12 May 2003** | **SCIA 5.04 to GOME WFD 12 May 2003**



Latitude dependence



Solar zenith angle (SZA) dependence



Total ozone dependence



SCIA 5.01/5.04 to GOME WFD for Jan to June 2003





Validation of SCIA tO_3 v5.01/5.04 (~ outdated GOME GDP 2.4) to half-year data set of GOME WFD:

- offset of -1%(±2%)
- SCIA V5 shows strong SZA, tO_3, meridional and seasonal dependence



Preliminary conclusion from comparisons of SCIA WFD to GOME WFD:

- agrees to within 1%
- no bias
- weak dependencies to SZA, tO_3, latitude

Comparisons of SCIAMACHY WFD to OMI on 11th Nov 2004

OMI-level-3product from http://toms.gsfc.nasa.gov/pub/omi | **binned SCIA WFD** | **binned GOME WFD** from http://www.iup.uni-bremen.de/gome/wfdoas | **OMI-level-3 product** from http://toms.gsfc.nasa.gov/pub/omi | **binned SCIA WFD**



Comparison between OMI-L3 and SCIA WFD (binned into 1° lat x 1.25° lon)

GOME WFD & SCIA WFD | **Overview** | **latitude dependence** | **SZA dependence** | **tO_3 dependence**



Conclusions & Outlook

Results from GOME WFD show excellent agreement with Brewer measurements.

The WFDOAS algorithm has been successfully adapted to SCIAMACHY and shows large improvements in comparison to recent operational SCIAMACHY 5.04 data with negligible dependencies to latitude, SZA and tO_3.

Preliminary comparisons of current OMI level-3 product and SCIAMACHY WFDOAS show small negative bias (≤2%) of OMI to SCIA WFD which is more pronounced at low tO_3 and SZA >60°.

Reprocessing of the entire SCIAMACHY data set with WFDOAS is underway and an extensive validation of this data set is planned.

Further comparisons between SCIAMACHY, GOME and OMI will clarify how these different data sets can be combined for tO_3 trend studies.

References & Acknowledgements

Bracher A., Lamsal L.N., Weber M., Bramstedt K., Coldewey-Egbers M., Burrows J. P. (2005) Global satellite validation of SCIAMACHY ozone columns with GOME WFDOAS. Atmos. Chem. Phys. 5: 2357-2368.
Coldewey-Egbers M., Weber M., Lamsal L.N., de Beek R., Buchwitz M., Burrows J.P. (2005) Total ozone retrieval from GOME UV spectral data using the weighting function DOAS approach. Atmos. Chem. Phys. 5: 1015-1025.
Weber M., Lamsal L.N., Coldewey-Egbers M., Bramstedt K., Burrows J. P. (2005) Pole-to-pole validation of GOME WFDOAS total ozone with groundbased data. Atmos. Chem. Phys. 5:1341-1355.

We thank DLR and ESA/ESRIN for providing GOME and SCIAMACHY calibrated level 1 spectral, and NASA and the OMI/AURA team for OMI level 3 data. This work is funded in part by ESA-ESRIN (projects SATVAL and SciLoV), the DLR-Bonn (contract No. 50 EE0025, 50 EE0502), the BMBF (FKZ 01 SF9994 and 7ATF42 (GOMSTRAT) within the AFO 2000 national research program), and EU project EVK2-CT-2001-00133 (CANDIDOZ). The SCIAMACHY and GOME WFDOAS data shown here were calculated on the HLRN (High-Performance Computer Center North). Services and support are gratefully acknowledged.

Global comparisons of total O3 columns from SCIAMACHY, GOME and OMI

A. Bracher, L. N. Lamsal, M. Weber, M. Coldewey-Egbers, J. P. Burrows
University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de

Introduction

Global stratospheric ozone columns derived from UV nadir spectra measured by OMI, SCIAMACHY and GOME are compared. Data sets for SCIAMACHY are the products from the weighting function DOAS algorithm (WFDOAS Version 1; see Coldewey-Egbers et al. 2005) and the most recent ESA operational product versions 5.04/5.01, GOME WFDOAS and OMI level-3 data are taken from the are from IUP webpage and TOMS website, respectively Because of different spatial resolutions (GOME 320x40, SCIAMACHY 60x30, OMI 24x18 km), data from 2003 and 2005 are binned into appropriate spatial bins and then compared.

GOME Weighting Function DOAS (WFDOAS): Algorithm and Validation

Main feature of WFDOAS (see Coldewey-Egbers et al. (2005):

- Taylor expansion of optical depth spectrum around climatological total ozone (TOZ_{clim})
- Modelled radiative transfer quantities: reference OD and weighting function parameterised by : TOZ, eff. albedo, eff. scene height, viewing geometry
- Improved Raman correction
- Ozone temperature correction
- Cloud cover and effective scene height from oxygen A-band absorption (Koellemeijer et al. 2001) including ghost vertical column correction
- Effective UV albedo (surface reflectance) from 377nm
- Spectral fitting window: 326.6-335nm
- Climatological ozone from TOMS V7 profile shape climatology (Wellemeyer et al. 1997)

Comparison of GOME WFDOAS with Brewer data (see Weber et al. 2005)

Well-maintained Brewer data- good source for satellite O3 validation since Brewer has negligible O3-T dependence in their wavelengths



GOME WFD compared with Brewer data: within 1% and negligible seasonal cycle!

Comparisons of GOME WFD, SCIAMACHY 5.01/5.04 and SCIAMACHY WFD (from Bracher et al. 2005)

SCIAMACHY (=SCIA) WFD V1, SCIA operational product 5.04 and GOME WFD V1 are compared at several days during Jan to June 2003, the comparison between SCIA op. 5.04 and GOME WFD V1 was performed for the whole data set during this time frame. GOME is a good candidate for validation of SCIA: equator crossing time 10:30 LT (SCIA 10 LT) and same orbit. Data sets are binned into 2.5° lat x 2.5° lon.

Global tO_3 distributions



Overview

SCIA WFD to GOME WFD 12 May 2003 | **SCIA WFD to SCIA 5.04 12 May 2003** | **SCIA 5.04 to GOME WFD 12 May 2003**



Latitude dependence



Solar zenith angle (SZA) dependence



Total ozone dependence



SCIA 5.01/5.04 to GOME WFD for Jan to June 2003





Validation of SCIA tO_3 v5.01/5.04 (~ outdated GOME GDP 2.4) to half-year data set of GOME WFD:

- offset of -1%(±2%)
- SCIA V5 shows strong SZA, tO_3, meridional and seasonal dependence



Preliminary conclusion from comparisons of SCIA WFD to GOME WFD:

- agrees to within 1%
- no bias
- weak dependencies to SZA, tO_3, latitude

Comparisons of SCIAMACHY WFD to OMI on 11th Nov 2004

OMI-level-3product from http://toms.gsfc.nasa.gov/pub/omi | **binned SCIA WFD** | **binned GOME WFD** from http://www.iup.uni-bremen.de/gome/wfdoas | **OMI-level-3 product** from http://toms.gsfc.nasa.gov/pub/omi | **binned SCIA WFD**



Comparison between OMI-L3 and SCIA WFD (binned into 1° lat x 1.25° lon)

GOME WFD & SCIA WFD | **Overview** | **latitude dependence** | **SZA dependence** | **tO_3 dependence**



Conclusions & Outlook

Results from GOME WFD show excellent agreement with Brewer measurements.

The WFDOAS algorithm has been successfully adapted to SCIAMACHY and shows large improvements in comparison to recent operational SCIAMACHY 5.04 data with negligible dependencies to latitude, SZA and tO_3.

Preliminary comparisons of current OMI level-3 product and SCIAMACHY WFDOAS show small negative bias (≤2%) of OMI to SCIA WFD which is more pronounced at low tO_3 and SZA >60°.

Reprocessing of the entire SCIAMACHY data set with WFDOAS is underway and an extensive validation of this data set is planned.

Further comparisons between SCIAMACHY, GOME and OMI will clarify how these different data sets can be combined for tO_3 trend studies.

References & Acknowledgements

Bracher A., Lamsal L.N., Weber M., Bramstedt K., Coldewey-Egbers M., Burrows J. P. (2005) Global satellite validation of SCIAMACHY ozone columns with GOME WFDOAS. Atmos. Chem. Phys. 5: 2357-2368.
Coldewey-Egbers M., Weber M., Lamsal L.N., de Beek R., Buchwitz M., Burrows J.P. (2005) Total ozone retrieval from GOME UV spectral data using the weighting function DOAS approach. Atmos. Chem. Phys. 5: 1015-1025.
Weber M., Lamsal L.N., Coldewey-Egbers M., Bramstedt K., Burrows J. P. (2005) Pole-to-pole validation of GOME WFDOAS total ozone with groundbased data. Atmos. Chem. Phys. 5:1341-1355.

We thank DLR and ESA/ESRIN for providing GOME and SCIAMACHY calibrated level 1 spectral, and NASA and the OMI/AURA team for OMI level 3 data. This work is funded in part by ESA-ESRIN (projects SATVAL and SciLoV), the DLR-Bonn (contract No. 50 EE0025, 50 EE0502), the BMBF (FKZ 01 SF9994 and 7ATF42 (GOMSTRAT) within the AFO 2000 national research program), and EU project EVK2-CT-2001-00133 (CANDIDOZ). The SCIAMACHY and GOME WFDOAS data shown here were calculated on the HLRN (High-Performance Computer Center North). Services and support are gratefully acknowledged.

Global comparisons of total O3 columns from SCIAMACHY, GOME and OMI

A. Bracher, L. N. Lamsal, M. Weber, M. Coldewey-Egbers, J. P. Burrows
University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de

Introduction

Global stratospheric ozone columns derived from UV nadir spectra measured by OMI, SCIAMACHY and GOME are compared. Data sets for SCIAMACHY are the products from the weighting function DOAS algorithm (WFDOAS Version 1; see Coldewey-Egbers et al. 2005) and the most recent ESA operational product versions 5.04/5.01, GOME WFDOAS and OMI level-3 data are taken from the are from IUP webpage and TOMS website, respectively Because of different spatial resolutions (GOME 320x40, SCIAMACHY 60x30, OMI 24x18 km), data from 2003 and 2005 are binned into appropriate spatial bins and then compared.

GOME Weighting Function DOAS (WFDOAS): Algorithm and Validation

Main feature of WFDOAS (see Coldewey-Egbers et al. (2005):

- Taylor expansion of optical depth spectrum around climatological total ozone (TOZ_{clim})
- Modelled radiative transfer quantities: reference OD and weighting function parameterised by : TOZ, eff. albedo, eff. scene height, viewing geometry
- Improved Raman correction
- Ozone temperature correction
- Cloud cover and effective scene height from oxygen A-band absorption (Koellemeijer et al. 2001) including ghost vertical column correction
- Effective UV albedo (surface reflectance) from 377nm
- Spectral fitting window: 326.6-335nm
- Climatological ozone from TOMS V7 profile shape climatology (Wellemeyer et al. 1997)

Comparison of GOME WFDOAS with Brewer data (see Weber et al. 2005)

Well-maintained Brewer data- good source for satellite O3 validation since Brewer has negligible O3-T dependence in their wavelengths



GOME WFD compared with Brewer data: within 1% and negligible seasonal cycle!

Comparisons of GOME WFD, SCIAMACHY 5.01/5.04 and SCIAMACHY WFD (from Bracher et al. 2005)

SCIAMACHY (=SCIA) WFD V1, SCIA operational product 5.04 and GOME WFD V1 are compared at several days during Jan to June 2003, the comparison between SCIA op. 5.04 and GOME WFD V1 was performed for the whole data set during this time frame. GOME is a good candidate for validation of SCIA: equator crossing time 10:30 LT (SCIA 10 LT) and same orbit. Data sets are binned into 2.5° lat x 2.5° lon.

Global tO_3 distributions



Overview

SCIA WFD to GOME WFD 12 May 2003 | **SCIA WFD to SCIA 5.04 12 May 2003** | **SCIA 5.04 to GOME WFD 12 May 2003**



Latitude dependence



Solar zenith angle (SZA) dependence



Total ozone dependence



SCIA 5.01/5.04 to GOME WFD for Jan to June 2003





Validation of SCIA tO_3 v5.01/5.04 (~ outdated GOME GDP 2.4) to half-year data set of GOME WFD:

- offset of -1%(±2%)
- SCIA V5 shows strong SZA, tO_3, meridional and seasonal dependence



Preliminary conclusion from comparisons of SCIA WFD to GOME WFD:

- agrees to within 1%
- no bias
- weak dependencies to SZA, tO_3, latitude

Comparisons of SCIAMACHY WFD to OMI on 11th Nov 2004

OMI-level-3product from http://toms.gsfc.nasa.gov/pub/omi | **binned SCIA WFD** | **binned GOME WFD** from http://www.iup.uni-bremen.de/gome/wfdoas | **OMI-level-3 product** from http://toms.gsfc.nasa.gov/pub/omi | **binned SCIA WFD**



Comparison between OMI-L3 and SCIA WFD (binned into 1° lat x 1.25° lon)

GOME WFD & SCIA WFD | **Overview** | **latitude dependence** | **SZA dependence** | **tO_3 dependence**



Conclusions & Outlook

Results from GOME WFD show excellent agreement with Brewer measurements.

The WFDOAS algorithm has been successfully adapted to SCIAMACHY and shows large improvements in comparison to recent operational SCIAMACHY 5.04 data with negligible dependencies to latitude, SZA and tO_3.

Preliminary comparisons of current OMI level-3 product and SCIAMACHY WFDOAS show small negative bias (≤2%) of OMI to SCIA WFD which is more pronounced at low tO_3 and SZA >60°.

Reprocessing of the entire SCIAMACHY data set with WFDOAS is underway and an extensive validation of this data set is planned.

Further comparisons between SCIAMACHY, GOME and OMI will clarify how these different data sets can be combined for tO_3 trend studies.

References & Acknowledgements

Bracher A., Lamsal L.N., Weber M., Bramstedt K., Coldewey-Egbers M., Burrows J. P. (2005) Global satellite validation of SCIAMACHY ozone columns with GOME WFDOAS. Atmos. Chem. Phys. 5: 2357-2368.
Coldewey-Egbers M., Weber M., Lamsal L.N., de Beek R., Buchwitz M., Burrows J.P. (2005) Total ozone retrieval from GOME UV spectral data using the weighting function DOAS approach. Atmos. Chem. Phys. 5: 1015-1025.
Weber M., Lamsal L.N., Coldewey-Egbers M., Bramstedt K., Burrows J. P. (2005) Pole-to-pole validation of GOME WFDOAS total ozone with groundbased data. Atmos. Chem. Phys. 5:1341-1355.

We thank DLR and ESA/ESRIN for providing GOME and SCIAMACHY calibrated level 1 spectral, and NASA and the OMI/AURA team for OMI level 3 data. This work is funded in part by ESA-ESRIN (projects SATVAL and SciLoV), the DLR-Bonn (contract No. 50 EE0025, 50 EE0502), the BMBF (FKZ 01 SF9994 and 7ATF42 (GOMSTRAT) within the AFO 2000 national research program), and EU project EVK2-CT-2001-00133 (CANDIDOZ). The SCIAMACHY and GOME WFDOAS data shown here were calculated on the HLRN (High-Performance Computer Center North). Services and support are gratefully acknowledged.

Global comparisons of total O3 columns from SCIAMACHY, GOME and OMI

A. Bracher, L. N. Lamsal, M. Weber, M. Coldewey-Egbers, J. P. Burrows
University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de

Introduction

Global stratospheric ozone columns derived from UV nadir spectra measured by OMI, SCIAMACHY and GOME are compared. Data sets for SCIAMACHY are the products from the weighting function DOAS algorithm (WFDOAS Version 1; see Coldewey-Egbers et al. 2005) and the most recent ESA operational product versions 5.04/5.01, GOME WFDOAS and OMI level-3 data are taken from the are from IUP webpage and TOMS website, respectively Because of different spatial resolutions (GOME 320x40, SCIAMACHY 60x30, OMI 24x18 km), data from 2003 and 2005 are binned into appropriate spatial bins and then compared.

GOME Weighting Function DOAS (WFDOAS): Algorithm and Validation

Main feature of WFDOAS (see Coldewey-Egbers et al. (2005):

- Taylor expansion of optical depth spectrum around climatological total ozone (TOZ_{clim})
- Modelled radiative transfer quantities: reference OD and weighting function parameterised by : TOZ, eff. albedo, eff. scene height, viewing geometry
- Improved Raman correction
- Ozone temperature correction
- Cloud cover and effective scene height from oxygen A-band absorption (Koellemeijer et al. 2001) including ghost vertical column correction
- Effective UV albedo (surface reflectance) from 377nm
- Spectral fitting window: 326.6-335nm
- Climatological ozone from TOMS V7 profile shape climatology (Wellemeyer et al. 1997)

Comparison of GOME WFDOAS with Brewer data (see Weber et al. 2005)

Well-maintained Brewer data- good source for satellite O3 validation since Brewer has negligible O3-T dependence in their wavelengths



GOME WFD compared with Brewer data: within 1% and negligible seasonal cycle!

Comparisons of GOME WFD, SCIAMACHY 5.01/5.04 and SCIAMACHY WFD (from Bracher et al. 2005)

SCIAMACHY (=SCIA) WFD V1, SCIA operational product 5.04 and GOME WFD V1 are compared at several days during Jan to June 2003, the comparison between SCIA op. 5.04 and GOME WFD V1 was performed for the whole data set during this time frame. GOME is a good candidate for validation of SCIA: equator crossing time 10:30 LT (SCIA 10 LT) and same orbit. Data sets are binned into 2.5° lat x 2.5° lon.

Global tO_3 distributions



Overview

SCIA WFD to GOME WFD 12 May 2003 | **SCIA WFD to SCIA 5.04 12 May 2003** | **SCIA 5.04 to GOME WFD 12 May 2003**



Latitude dependence



Solar zenith angle (SZA) dependence



Total ozone dependence



SCIA 5.01/5.04 to GOME WFD for Jan to June 2003





Validation of SCIA tO_3 v5.01/5.04 (~ outdated GOME GDP 2.4) to half-year data set of GOME WFD:

- offset of -1%(±2%)
- SCIA V5 shows strong SZA, tO_3, meridional and seasonal dependence



Preliminary conclusion from comparisons of SCIA WFD to GOME WFD:

- agrees to within 1%
- no bias
- weak dependencies to SZA, tO_3, latitude

Comparisons of SCIAMACHY WFD to OMI on 11th Nov 2004

OMI-level-3product from http://toms.gsfc.nasa.gov/pub/omi | **binned SCIA WFD** | **binned GOME WFD** from http://www.iup.uni-bremen.de/gome/wfdoas | **OMI-level-3 product** from http://toms.gsfc.nasa.gov/pub/omi | **binned SCIA WFD**



Comparison between OMI-L3 and SCIA WFD (binned into 1° lat x 1.25° lon)

GOME WFD & SCIA WFD | **Overview** | **latitude dependence** | **SZA dependence** | **tO_3 dependence**



Conclusions & Outlook

Results from GOME WFD show excellent agreement with Brewer measurements.

The WFDOAS algorithm has been successfully adapted to SCIAMACHY and shows large improvements in comparison to recent operational SCIAMACHY 5.04 data with negligible dependencies to latitude, SZA and tO_3.

Preliminary comparisons of current OMI level-3 product and SCIAMACHY WFDOAS show small negative bias (≤2%) of OMI to SCIA WFD which is more pronounced at low tO_3 and SZA >60°.

Reprocessing of the entire SCIAMACHY data set with WFDOAS is underway and an extensive validation of this data set is planned.

Further comparisons between SCIAMACHY, GOME and OMI will clarify how these different data sets can be combined for tO_3 trend studies.

References & Acknowledgements

Bracher A., Lamsal L.N., Weber M., Bramstedt K., Coldewey-Egbers M., Burrows J. P. (2005) Global satellite validation of SCIAMACHY ozone columns with GOME WFDOAS. Atmos. Chem. Phys. 5: 2357-2368.
Coldewey-Egbers M., Weber M., Lamsal L.N., de Beek R., Buchwitz M., Burrows J.P. (2005) Total ozone retrieval from GOME UV spectral data using the weighting function DOAS approach. Atmos. Chem. Phys. 5: 1015-1025.
Weber M., Lamsal L.N., Coldewey-Egbers M., Bramstedt K., Burrows J. P. (2005) Pole-to-pole validation of GOME WFDOAS total ozone with groundbased data. Atmos. Chem. Phys. 5:1341-1355.

We thank DLR and ESA/ESRIN for providing GOME and SCIAMACHY calibrated level 1 spectral, and NASA and the OMI/AURA team for OMI level 3 data. This work is funded in part by ESA-ESRIN (projects SATVAL and SciLoV), the DLR-Bonn (contract No. 50 EE0025, 50 EE0502), the BMBF (FKZ 01 SF9994 and 7ATF42 (GOMSTRAT) within the AFO 2000 national research program), and EU project EVK2-CT-2001-00133 (CANDIDOZ). The SCIAMACHY and GOME WFDOAS data shown here were calculated on the HLRN (High-Performance Computer Center North). Services and support are gratefully acknowledged.

Global comparisons of total O3 columns from SCIAMACHY, GOME and OMI

A. Bracher, L. N. Lamsal, M. Weber, M. Coldewey-Egbers, J. P. Burrows
University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de

Introduction

Global stratospheric ozone columns derived from UV nadir spectra measured by OMI, SCIAMACHY and GOME are compared. Data sets for SCIAMACHY are the products from the weighting function DOAS algorithm (WFDOAS Version 1; see Coldewey-Egbers et al. 2005) and the most recent ESA operational product versions 5.04/5.01, GOME WFDOAS and OMI level-3 data are taken from the are from IUP webpage and TOMS website, respectively Because of different spatial resolutions (GOME 320x40, SCIAMACHY 60x30, OMI 24x18 km), data from 2003 and 2005 are binned into appropriate spatial bins and then compared.

GOME Weighting Function DOAS (WFDOAS): Algorithm and Validation

Main feature of WFDOAS (see Coldewey-Egbers et al. (2005):

- Taylor expansion of optical depth spectrum around climatological total ozone (TOZ_{clim})
- Modelled radiative transfer quantities: reference OD and weighting function parameterised by : TOZ, eff. albedo, eff. scene height, viewing geometry
- Improved Raman correction
- Ozone temperature correction
- Cloud cover and effective scene height from oxygen A-band absorption (Koellemeijer et al. 2001) including ghost vertical column correction
- Effective UV albedo (surface reflectance) from 377nm
- Spectral fitting window: 326.6-335nm
- Climatological ozone from TOMS V7 profile shape climatology (Wellemeyer et al. 1997)

Comparison of GOME WFDOAS with Brewer data (see Weber et al. 2005)

Well-maintained Brewer data- good source for satellite O3 validation since Brewer has negligible O3-T dependence in their wavelengths



GOME WFD compared with Brewer data: within 1% and negligible seasonal cycle!

Comparisons of GOME WFD, SCIAMACHY 5.01/5.04 and SCIAMACHY WFD (from Bracher et al. 2005)

SCIAMACHY (=SCIA) WFD V1, SCIA operational product 5.04 and GOME WFD V1 are compared at several days during Jan to June 2003, the comparison between SCIA op. 5.04 and GOME WFD V1 was performed for the whole data set during this time frame. GOME is a good candidate for validation of SCIA: equator crossing time 10:30 LT (SCIA 10 LT) and same orbit. Data sets are binned into 2.5° lat x 2.5° lon.

Global tO_3 distributions



Overview

SCIA WFD to GOME WFD 12 May 2003 | **SCIA WFD to SCIA 5.04 12 May 2003** | **SCIA 5.04 to GOME WFD 12 May 2003**



Latitude dependence



Solar zenith angle (SZA) dependence



Total ozone dependence



SCIA 5.01/5.04 to GOME WFD for Jan to June 2003





Validation of SCIA tO_3 v5.01/5.04 (~ outdated GOME GDP 2.4) to half-year data set of GOME WFD:

- offset of -1%(±2%)
- SCIA V5 shows strong SZA, tO_3, meridional and seasonal dependence



Preliminary conclusion from comparisons of SCIA WFD to GOME WFD:

- agrees to within 1%
- no bias
- weak dependencies to SZA, tO_3, latitude

Comparisons of SCIAMACHY WFD to OMI on 11th Nov 2004

OMI-level-3product from http://toms.gsfc.nasa.gov/pub/omi | **binned SCIA WFD** | **binned GOME WFD** from http://www.iup.uni-bremen.de/gome/wfdoas | **OMI-level-3 product** from http://toms.gsfc.nasa.gov/pub/omi | **binned SCIA WFD**



Comparison between OMI-L3 and SCIA WFD (binned into 1° lat x 1.25° lon)

GOME WFD & SCIA WFD | **Overview** | **latitude dependence** | **SZA dependence** | **tO_3 dependence**



Conclusions & Outlook

Results from GOME WFD show excellent agreement with Brewer measurements.

The WFDOAS algorithm has been successfully adapted to SCIAMACHY and shows large improvements in comparison to recent operational SCIAMACHY 5.04 data with negligible dependencies to latitude, SZA and tO_3.

Preliminary comparisons of current OMI level-3 product and SCIAMACHY WFDOAS show small negative bias (≤2%) of OMI to SCIA WFD which is more pronounced at low tO_3 and SZA >60°.

Reprocessing of the entire SCIAMACHY data set with WFDOAS is underway and an extensive validation of this data set is planned.

Further comparisons between SCIAMACHY, GOME and OMI will clarify how these different data sets can be combined for tO_3 trend studies.

References & Acknowledgements

Bracher A., Lamsal L.N., Weber M., Bramstedt K., Coldewey-Egbers M., Burrows J. P. (2005) Global satellite validation of SCIAMACHY ozone columns with GOME WFDOAS. Atmos. Chem. Phys. 5: 2357-2368.
Coldewey-Egbers M., Weber M., Lamsal L.N., de Beek R., Buchwitz M., Burrows J.P. (2005) Total ozone retrieval from GOME UV spectral data using the weighting function DOAS approach. Atmos. Chem. Phys. 5: 1015-1025.
Weber M., Lamsal L.N., Coldewey-Egbers M., Bramstedt K., Burrows J. P. (2005) Pole-to-pole validation of GOME WFDOAS total ozone with groundbased data. Atmos. Chem. Phys. 5:1341-1355.

We thank DLR and ESA/ESRIN for providing GOME and SCIAMACHY calibrated level 1 spectral, and NASA and the OMI/AURA team for OMI level 3 data. This work is funded in part by ESA-ESRIN (projects SATVAL and SciLoV), the DLR-Bonn (contract No. 50 EE0025, 50 EE0502), the BMBF (FKZ 01 SF9994 and 7ATF42 (GOMSTRAT) within the AFO 2000 national research program), and EU project EVK2-CT-2001-00133 (CANDIDOZ). The SCIAMACHY and GOME WFDOAS data shown here were calculated on the HLRN (High-Performance Computer Center North). Services and support are gratefully acknowledged.